arXiv:1412.6330v1 [math.DG] 19 Dec 2014

ON LORENTZIAN TWO-SYMMETRIC MANIFOLDS OF DIMENSION FOUR

A. ZAEIM, M. CHAICHI AND Y. KESHAVARZI

Abstract. We study curvature properties of four-dimensional Lorentzian manifold with two-symmetry property. We then consider Einstein-like metrics, Ricci solitons and homogeneity over these spaces.

1. Introduction

Symmetric manifolds are a large class of spaces with many interesting geometric properties. A direct generalization of these spaces could be realized by so-called k-symmetry property. A (pseudo-)Riemannian space (M, g) is called k-symmetric if the following condition is valid:

$$\nabla^k R = 0, \quad \nabla^{k-1} R \neq 0,$$

where $k \geq 1$ and R is the curvature tensor of (M, g). In the Riemannian setting, contrary to the pseudo-Riemannian case, a k-symmetric space is necessarily locally symmetric, i.e., $\nabla R = 0$ [23]. Examples of pseudo-Riemannian k-symmetric spaces with $k \geq 2$ can be found in [22, 6, 18]. Many interesting results about Lorentzian two-symmetric spaces were presented in [22], in particular the author proved that any two-symmetric Lorentzian manifold admits a parallel null vector field. A classification of four-dimensional two-symmetric Lorentzian spaces is obtained in [6], and it is shown that such spaces are some special pp-waves. This result is based on the Petrov classification of the Weyl tensors. The generalization of the results of [6] is the subject of [3], where it is proven that a locally indecomposable Lorentzian manifold of dimension $n + 2$ is two-symmetric if and only if there exist local coordinates $(v, x^1, \ldots, x^n, u)$ such that

$$g = 2dvdu + \sum_{i=1}^{n} (dx^i)^2 + (H_{ij}u + F_{ij})x^i x^j (du)^2, \tag{1.1}$$

where H_{ij} is a nonzero diagonal matrix with diagonal elements $\lambda_1 \leq \cdots \leq \lambda_n$, and F_{ij} is a symmetric real matrix. According to this general form of Lorentzian two-symmetric manifolds, in the four-dimensional case, there exist local coordinates $(x^1, \ldots, x^4)$ such that the metric g of a Lorentzian two-symmetric space is

$$g = 2dx^1dx^4 + (dx^2)^2 + (dx^3)^2 + \left(x^4(a(x^2)^2 + b(x^3)^2) + p(x^2)^2 + 2qx^2x^3 + s(x^3)^2\right)(dx^4)^2, \tag{1.2}$$

2010 *Mathematics Subject Classification.* 53B30, 53C35, 53C30.

Key words and phrases. Pseudo-Riemannian metric, Einsteil-like metrics, Ricci soliton, Homogeneous space.

where a, b, p, q, s are real constants and $a^2 + b^2 \neq 0$. Our main goal is to study some geometric properties of four-dimensional Lorentzian two-symmetric spaces.

This paper is organized in the following way. Curvature properties of Lorentzian two-symmetric four-spaces will be studied in the section two and Einstein-like metrics of the spaces under consideration is the subject of section three. Ricci solitons and homogeneous four-dimensional Lorentzian two-symmetric spaces will be studied in section four and five respectively.

2. Two-symmetric Lorentzian four-manifolds

The first step for study the geometry of (pseudo-)Riemannian manifolds is to determine the Lievi-Civita connection. By using the *Koszul* identity $2g(\nabla_X Y, Z) = Xg(Y, Z) + Yg(Z, X) - Zg(X, Y) - g(X, [Y, Z]) + g(Y, [Z, X]) + g(Z, [X, Y])$, and applying the metric (1.2), one can determine the components of the Levi-Civita connection. We use $\partial_i = \frac{\partial}{\partial x^i}$ as a local basis for the tangent space and have:

Theorem 2.1. *Let (M, g) be an arbitrary two-symmetric Lorentzian four-manifold, where the metric g is described in local coordinates (x^1, x^2, x^3, x^4) by the Equation* (1.2)*. The non-zero components of the Levi-Civita connection are:*

$$\begin{aligned}
&\nabla_{\partial_2}\partial_4 = (ax^2x^4 + px^2 + qx^3)\partial_1,\\
&\nabla_{\partial_3}\partial_4 = (bx^3x^4 + sx^3 + qx^2)\partial_1,\\
&\nabla_{\partial_4}\partial_4 = \tfrac{a(x^2)^2+b(x^3)^2}{2}\partial_1 - (ax^2x^4 + px^2 + qx^3)\partial_2 - (bx^3x^4 + qx^2 + sx^3)\partial_3.
\end{aligned} \tag{2.3}$$

Applying the relation $R(X, Y) = [\nabla_X, \nabla_Y] - \nabla_{[X,Y]}$ we immediately determine the curvature tensor. If we set $R(\partial_k, \partial_l)\partial_j = R^i_{jkl}\partial_i$, then by contraction on the first and third indices of the curvature tensor, the Ricci tensor ϱ will be deduced. The scalar curvature tensor τ is also obtained by full contraction of coefficients of the curvature tensor.

Theorem 2.2. *A four-dimensional two-symmetric Lorentzian space admits zero scalar curvature. Also, the non-zero components of curvature tensor and Ricci tensor are*

$$\begin{aligned}
&R(\partial_2, \partial_4) = (ax^4 + p)\partial_1 dx^2 + q\partial_1 dx^3 - (ax^4 + p)\partial_2 dx^4 - q\partial_3 dx^4,\\
&R(\partial_3, \partial_4) = q\partial_1 dx^2 + (bx^4 + s)\partial_1 dx^3 - q\partial_2 dx^4 - (bx^4 + s)\partial_3 dx^4,\\
&\varrho(\partial_4, \partial_4) = -(a + b)x^4 - (s + p).
\end{aligned} \tag{2.4}$$

A (pseudo-)Riemannian manifold (M, g) is called *Einstein* if $\varrho = cg$, for a real constant c. Being *Ricci flat* means that the Ricci tensor vanishes identically. Also, *conformal flatness* translates into the following system of algebraic equations:

$$W_{ijkh} = R_{ijkh} - \tfrac{1}{2}(g_{ik}\varrho_{jh} + g_{jh}\varrho_{ik} - g_{ih}\varrho_{jk} - g_{jk}\varrho_{ih}) + \tfrac{r}{6}(g_{ik}g_{jh} - g_{ih}g_{jk}) = 0 \tag{2.5}$$

for all indices $i, j, k, h = 1, \ldots, 4$,

where W denotes the *Weyl tensor* and τ is the scalar curvature. Although two-symmetric spaces clearly aren't flat, we can check Ricci flatness.

Theorem 2.3. *Let (M, g) be a two-symmetric four-dimensional Lorentzian manifold such that the metric g is described by the Equation (1.2) in local coordinates (x^1, x^2, x^3, x^4). The following statements satisfy:*

a) *(M, g) is Einstein if and only if be Ricci flat if and only if $b = -a,\ s = -p$.*
b) *(M, g) is conformally flat if and only if $b = a,\ s = p,\ q = 0$.*

Proof. Let (M, g) be an Einstein manifold. Using the Equation (2.4) we set $\varrho = cg$. The following relations must be established.

$$c = (a + b)x^4 + s + p = 0.$$

So the Einstein property is equivalent to satisfying $b = -a$, $s = -p$ and $c = 0$ which is clearly equivalent to Ricci flatness. Using the Equation (2.5), the non-zero components of the Weyl tensor would be

$$W_{2424} = -W_{3434} = \tfrac{(b-a)x^4+s-p}{2}, \quad W_{2434} = -q,$$

it is obvious that the Weyl tensor vanishes if and only if $b = a$, $s = p$ and $q = 0$. □

3. Einstein-like Lorentzian two-symmetric spaces

Two new classes of Riemannian manifolds which are defined through conditions on the Ricci tensor, introduced by A. Gray in [17]. These types of manifolds which are famous as $\mathcal{A}$ and $\mathcal{B}$ classes, would be extended at once to the pseudo-Riemannian geometry. $\mathcal{A}$ and $\mathcal{B}$ classes are defined in the following way:

Class $\mathcal{A}$: a pseudo-Riemannian manifold (M, g) belongs to class $\mathcal{A}$ if and only if its Ricci tensor ϱ is *cyclic-parallel*, that is,

$$(\nabla_X \varrho)(Y, Z) + (\nabla_Y \varrho)(Z, X) + (\nabla_Z \varrho)(X, Y) = 0, \tag{3.6}$$

for all vector fields X, Y and Z on M. The Equation (3.6) is equivalent to requiring that ϱ is a *Killing tensor*, that is,

$$(\nabla_X \varrho)(X, X) = 0. \tag{3.7}$$

To note that Equation (3.7), also known as the first odd Ledger condition, is a necessary condition for a (pseudo-)Riemannian manifold to be a D'Atri space. Hence, identifying two-symmetric manifolds of a given dimension satisfying (3.7), is the first step to understand D'Atri spaces.

Class $\mathcal{B}$: a pseudo-Riemannin manifold (M, g) belongs to class $\mathcal{B}$ if and only if its Ricci tensor be a *Codazzi tensor*, that is,

$$(\nabla_X \varrho)(Y, Z) = (\nabla_Y \varrho)(X, Z). \tag{3.8}$$

A pseudo-Riemnnain manifold which belongs to one of the above classes is called an *Einstein-like manifold*. We denote the class of Ricci parallel, Einstein and manifolds with constant scalar curvature by $\mathcal{P}$, $\mathcal{E}$ and $\mathcal{C}$ respectively. One can easily see that the intersection of two Einstein-like classes consists of Ricci parallel manifolds. This situation can be summarized in the following diagram:

$$\mathcal{E} \subset \mathcal{P} = \mathcal{A} \cap \mathcal{B} \begin{array}{l} \subset \mathcal{A} \\ \\ \subset \mathcal{B} \end{array} \subset \mathcal{A} \cup \mathcal{B} \subset \mathcal{C}.$$

Einstein-like metrics are deeply investigated through the different kinds of homogeneous spaces in both Riemannian and pseudo-Riemannian signatures. Three-dimensional Riemannian homogeneous spaces studied in [1]. In [8], authors study three- and four-dimensional Einstein-like Riemannian manifolds which are Ricci-curvature homogeneous. They could completely classify three-dimensional case of the mentioned spaces, while in the four-dimensional case, they partially classified the special case where the manifold is locally homogeneous. They also presented explicit examples of four-dimensional locally homogeneous Riemannian manifolds whose Ricci tensor is cyclic-parallel and has distinct eigenvalues. These examples invalidated the expectation stated by F. Podestá and A. Spiro in [21]. Three-dimensional ball-homogeneous spaces, semi-symmetric spaces, Sasakian spaces and three-dimensional contact metric manifolds are other Riemannian classes which were the subject of research for the Einstein-like properties [10, 7, 2, 11].

Theorem 3.1. *Every four-dimensional two-symmetric Lorentzian manifold (M, g) belongs to class $\mathcal{A}$, if and only if $b = -a$.*

Proof. Let $v = v^1\partial_1 + v^2\partial_2 + v^3\partial_3 + v^4\partial_4$ be an arbitrary vector space on (M, g), where $v^1, \dots, v^4$ are smooth functions on M. As mentioned before, (M, g) belongs to class $\mathcal{A}$ of Einstein-like manifolds if and only if the Equation (3.7) satisfies. By straight forward calculations it is implied that

$$(\nabla_v \varrho)(v, v) = -(v^4)^3(a + b).$$

Thus, (M, g) belongs to class $\mathcal{A}$ if and only if $b = -a$. □

Theorem 3.2. *Every four-dimensional two-symmetric Lorentzian manifold (M, g) belongs to class $\mathcal{B}$ of the Einstein-like manifolds.*

Proof. Let $v = \sum_{i=1}^4 v^i\partial_i$, $u = \sum_{i=1}^4 u^i\partial_i$ and $w = \sum_{i=1}^4 w^i\partial_i$ be three arbitrary smooth vector fields on (M, g). Every two-symmetric space (M, g) belongs to class $\mathcal{B}$ of Einstein-like manifolds if and only if the Equation (3.8) satisfies. Direct calculations yield that

$$(\nabla_u \varrho)(v, w) = (\nabla_v \varrho)(u, w) = -u^4v^4w^4(a + b),$$

and so, the Equation (3.8) always establishes and proves the claim. □

4. Two-symmetric Lorentzian Ricci solitons

We now report some basic information on Ricci solitons, referring to [14] for a survey and further references. A *Ricci soliton* is a pseudo-Riemannian manifold (M, g) admitting a smooth vector field V, such that

$$\mathcal{L}_V g + \varrho = \lambda g, \tag{4.9}$$

where $\mathcal{L}$ denotes the Lie derivative and λ a real constant. A Ricci soliton is said to be *shrinking*, *steady* or *expanding* depending on whether $\lambda > 0$, $\lambda = 0$ or $\lambda < 0$, respectively. Ricci solitons are self-similar solutions of the *Ricci flow*.

Originally introduced in the Riemannian case, Ricci solitons have been intensively studied in pseudo-Riemannian settings in recent years. The Ricci soliton equation is also a special case of Einstein field equations.

Theorem 4.1. *Every four-dimensional two-symmetric Lorentzian manifold (M, g) is shrinking, expanding and steady Ricci soliton.*

Proof. Let (M, g) be a four-dimensional two-symmetric Lorentzian manifold, where g is described by the Equation (1.2). Suppose that $v = \sum_{i=1}^{4} v^i \partial_i$ is a smooth vector field on (M, g) such that the Equation (4.9) satisfies for a real constant λ. The Lie derivative of g in direction v is

$$\begin{aligned}\mathcal{L}_v g = \; & 2\partial_1 v^4 (dx^1)^2 + 2(\partial_2 v^4 + \partial_1 v^2) dx^1 dx^2 + 2(\partial_3 v^4 + \partial_1 v^3) dx^1 dx^3 \\ & + 2(\partial_1 x^1 + \partial_4 v^4 + \partial_1 v^4 (a(x^2)^2 x^4 + p(x^2)^2 + 2qx^2 x^3 + b(x^3)^2 x^4 + s(x^3)^2)) dx^1 dx^4 + 2\partial_2 v^2 (dx^2)^2 + 2(\partial_2 v^3 + \partial_3 v^2) dx^2 dx^3 + 2(\partial_2 v^4 (a(x^2)^2 x^4 + \\ & p(x^2)^2 + 2qx^2 x^3 + b(x^3)^2 x^4 + s(x^3)^2 + 1) + \partial_2 v^1) dx^2 dx^4 + 2\partial_3 v^3 (dx^3)^2 + \\ & 2(\partial_3 v^4 (a(x^2)^2 x^4 + p(x^2)^2 + 2qx^2 x^3 + b(x^3)^2 x^4 + s(x^3)^2 + 1) + \partial_3 v^1) dx^3 dx^4 + \\ & (\partial_4 v^4 (2a(x^2)^2 x^4 + 2p(x^2)^2 + 4qx^2 x^3 + 2b(x^3)^2 (x^4) + 2s(x^3)^2) + 2\partial_4 v^1 + \\ & 2av^2 x^2 x^4 + 2pv^2 x^2 + 2qv^2 x^3 + 2qv^3 x^2 + 2bv^3 x^3 x^4 + 2sv^3 x^3 + av^4 (x^2)^2 + \\ & v^4 b(x^3)^2)(dx^4)^2.\end{aligned}$$

Applying Equations (1.2) and (2.4) in the Ricci soliton Equation (4.9), we have a system of PDEs which admits the following solution

$$\begin{cases} \lambda = 2c, \\ v^1 = \frac{(x^4)^2}{4}(a+b) + \frac{x^4}{2}(s+p) + 2cx^1, \\ v^2 = cx^2, \\ v^3 = cx^3, \\ v^4 = 0, \end{cases}$$

for a real constant c. Since c is arbitrary, (M, g) can be an expanding, shrinking or steady Ricci soliton. □

A Ricci soliton (M, g) is called *gradient* if the Equation (4.9) holds for a vector field $X = \mathrm{grad} f$, for some potential function f. In this case, (4.9) can be rewritten as

$$2\mathrm{Hes}_f + \varrho = \lambda g,$$

where Hes_f denotes the Hessian of f.

Theorem 4.2. *Every four-dimensional two-symmetric Lorentzian space (M, g) is a gradient Ricci soliton if and only if be a steady Ricci soliton. In this case, the potential function is $f = \frac{a+b}{12}(x^4)^3 + \frac{p+s}{4}(x^4)^2 + c_1 x^4 + c_2$, for arbitrary real constants c_1, c_2.*

Proof. Let $f = f(x^1, x^2, x^3, x^4)$ be a smooth function on (M, g) and $v = \sum_{i=1}^{4} v^i \partial_i$ be a gradient Ricci soliton with the potential function f. So the coefficient v^i must be $v^i = \sum_{j=1}^{4} g^{ij} \partial_j(f)$. By applying v to the Equation (4.9) the following equations must establish

$$\begin{cases} f_{11} = f_{12} = f_{13} = f_{23} = 0, \\ \lambda = 2f_{14} = 2f_{22} = 2f_{33}, \\ 2f_{24} - 2af_1 x^2 x^4 - 2pf_1 x^2 - 2qf_1 x^3 = 0, \\ 2f_{34} - 2bf_1 x^3 x^4 - 2qf_1 x^2 - 2sf_1 x^3 = 0, \\ \lambda(x^2)^2(ax^4 + p) + 2\lambda qx^2 x^3 + \lambda(x^3)^2(bx^4 + s) + (a+b)x^4 + s + p - 2f_{44} \\ + f_1(a(x^2)^2 + b(x^3)^2) - 2f_2(ax^2 x^4 + px^2 + qx^3) - 2f_3(qx^2 + bx^3 x^4 + sx^3) = 0, \end{cases}$$

where $f_i := \partial_i f$. After solving the above system of PDEs we get that λ must be vanished and f must be the same function of the statement, this matter ends the proof. □

5. Homogeneous two-symmetric four-dimentional spaces

Study of homogeneous spaces is one of the most interesting topics in differential geometry, where a deep connection between geometry and algebra appears. A (pseudo-)Riemannian manifold M is called *homogeneous*, if for any points $p, q \in M$, there is an isometry ϕ of M such that $\phi(p) = q$. In short, $I(M)$, the group of isometries of M, acts transitively on M. If M is homogeneous, then evidently any geometrical properties at one point of M holds at every point. For a detailed introduction to homogeneous spaces see e.g., [5, 20, 25].

Homogeneous Riemannian structures introduced by Ambrose and Singer in [4] and deeply studied in [24]. The notation is generalized to homogeneous pseudo-Riemannian structures by Gadea and Oubiña in [16], in order to obtain a characterization of reductive homogeneous pseudo-Riemannian manifolds. A pretty application of homogeneous structures on three dimensional Lorentzian manifold is shown in [9].

Let (M, g) be a connected pseudo-Riemannian manifold, the following definition introduced by Gadea and Oubiña:

Definition 5.1. [16] *A homogeneous pseudo-Riemannian structure on (M, g) is a tensor field T of type $(1, 2)$ on M, such that the connection $\widetilde{\nabla} = \nabla - T$ satisfies*

$$\widetilde{\nabla} g = 0, \quad \widetilde{\nabla} R = 0, \quad \widetilde{\nabla} T = 0. \tag{5.10}$$

The above conditions are equivalent to the following system of equations which are famous as Ambrose-Singer equations.

$$g(T_X Y, Z) + g(Y, T_X Z) = 0, \tag{5.11}$$

$$(\nabla_X R)_{YZ} = [T_X, R_{YZ}] - R_{T_X YZ} - R_{YT_X Z}, \tag{5.12}$$

$$(\nabla_X T)_Y = [T_X, T_Y] - T_{T_X Y}, \tag{5.13}$$

for all vector fields X, Y, Z.

Existence of homogeneous pseudo-Riemannian structures shows the homogeneity of the space. This fact is the subject of the following Lemma

Lemma 5.2. [16] *Let (M, g) be a simply connected and complete pseudo-Riemannian manifold, then (M, g) admits a pseudo-Riemannian homogeneous structure if and only if it is a reductive homogeneous pseudo-Riemannian manifold.*

Case1: Reductive cases:

By applying the above lemma, we consider reductive homogeneous four-dimensional two-symmetric Lorentzian spaces. The result is the following theorem,

Theorem 5.3. *Every Lorentzian four-dimensional two-symmetric manifold is not reductive homogeneous.*

Proof. Let (M, g) be a four-dimensional two-symmetric manifold. There exist local coordinates $(x^1, \ldots, x^4)$ such that the metric g is defined using the Equations (1.2). According to the Lemma 5.2, (M, g) is (locally) reductive homogeneous if and only if

the tensor field T of type $(1,2)$ exists, such that the Ambrose-Singer equations satisfy. Let $T_{\partial_i}\partial_j = T_{ij}^k\partial_k, 1 \leq i,j,k \leq 4$ be a homogeneous structure on (M,g) where $\partial_1 = \partial_v, \partial_2 = \partial_{x_1}, \partial_3 = \partial_{x_2}, \partial_4 = \partial_u$ and T_{ij}^k are smooth functions on M.
From the Equations (5.11) and (5.12), besides the relations between the components T_{ij}^k, one of the following relations for the constants a,b,p,q,s must be valid:

$$\begin{aligned}
&\mathbf{1}: a = p = q = 0,\\
&\mathbf{2}: a \neq 0, q = 0, s = \tfrac{bp}{a},\\
&\mathbf{3}: b = a, s = p, q = 0,\\
&\mathbf{4}: b = -a, s = -p,
\end{aligned}$$

but each of these solutions makes a contradiction with Equation (5.13) (or equivalently with $\widetilde{\nabla}T = 0$). For example, in the case 1, for the components T_{ij}^k have

$$\begin{aligned}
&T_{1j}^k = 0,\ 1 \leq j,k \leq 4, (j,k) \neq (4,1),\ T_{14}^1 = -T_{44}^4 = \tfrac{b}{2(bx^4+s)},\\
&T_{2j}^k = 0,\ 1 \leq j,k \leq 4,\ k \neq 1,\\
&T_{3j}^k = 0,\ 1 \leq j,k \leq 4,\ k \neq 1,\\
&T_{4j}^1 = 0,\ 1 \leq j \leq 3,\ T_{44}^1 = \tfrac{b(x^3)^2}{2},\\
&T_{4j}^k = -T_{kj}^1,\ 1 \leq j \leq 4,\ 2 \leq k \leq 3,\\
&T_{4j}^4 = 0,\ 1 \leq j \leq 3.
\end{aligned}$$

On the other hand, we have

$$\begin{aligned}
(\widetilde{\nabla}_{\partial_4}T)_{44}^4 = \ & \partial_4 T_{44}^4 + T_{42}^4(T_{44}^2 + x^2(p+ax^4) + qx^3) + T_{43}^4(T_{44}^3 + x^3(s+bx^4) + qx^2)\\
&+T_{24}^4(x^2(p+ax^4) + qx^3) + T_{34}^4(x^3(s+bx^4) + qx^2)\\
&+T_{41}^4(T_{44}^1 - \tfrac{a(x^2)^2+b(x^3)^2}{2}) - T_{14}^4\tfrac{a(x^2)^2+b(x^3)^2}{2} + (T_{44}^4)^2.
\end{aligned}$$

If we substitute the previous solutions in the above relation we get $(\widetilde{\nabla}_{\partial_4}T)_{44}^4 = \frac{3b^2}{4(bx^4+s)^2}$, and so the Equation (5.13) satisfies if $b = 0$ which is a contradiction, since in this case the matrix H in (1.1) vanishes. □

Case2: Non-reductive cases:

Consider a homogeneous manifold $M = G/H$ (with H connected), the Lie algebra $\mathfrak{g}$ of G , the isotropy subalgebra $\mathfrak{h}$, and $\mathfrak{m} = \mathfrak{g}/\mathfrak{h}$ the factor space, which identifies with a subspace of $\mathfrak{g}$ complementary to $\mathfrak{h}$. The pair $(\mathfrak{g},\mathfrak{h})$ uniquely defines the isotropy representation

$$\psi : \mathfrak{g} \to \mathfrak{gl}(\mathfrak{m}), \qquad \psi(x)(y) = [x,y]_{\mathfrak{m}} \quad \text{for all} \quad x \in \mathfrak{g}, y \in \mathfrak{m}.$$

Given a basis $\{h_1,...,h_r,u_1,...,u_n\}$ of $\mathfrak{g}$, where $\{h_j\}$ and $\{u_i\}$ are bases of $\mathfrak{h}$ and $\mathfrak{m}$, respectively, a bilinear form on $\mathfrak{m}$ is determined by the matrix g of its components with respect to the basis $\{u_i\}$ and is invariant if and only if ${}^t\psi(x)\circ g + g\circ\psi(x) = 0$ for all $x \in \mathfrak{h}$. Invariant pseudo-Riemannian metrics g on the homogeneous space $M = G/H$ are in a one-to-one correspondence with nondegenerate invariant symmetric bilinear forms g on $\mathfrak{m}$ [19].

Then, g uniquely defines its invariant linear Levi-Civita connection, described in terms of the corresponding homomorphism of $\mathfrak{h}$-modules $\Lambda : \mathfrak{g} \to \mathfrak{gl}(\mathfrak{m})$, such that $\Lambda(x)(y_{\mathfrak{m}}) = [x,y]_{\mathfrak{m}}$ for all $x \in \mathfrak{h}, y \in \mathfrak{g}$. Explicitly, one has

$$\Lambda(x)(y_{\mathfrak{m}}) = \tfrac{1}{2}[x,y]_{\mathfrak{m}} + v(x,y), \qquad \text{for all } x,y \in \mathfrak{g},$$

where $v : \mathfrak{g} \times \mathfrak{g} \to \mathfrak{m}$ is the $\mathfrak{h}$-invariant symmetric mapping uniquely determined by

$$2g(v(x,y),z_{\mathfrak{m}}) = g(x_{\mathfrak{m}},[z,y]_{\mathfrak{m}}) + g(y_{\mathfrak{m}},[z,x]_{\mathfrak{m}}), \qquad \text{for all } x,y,z \in \mathfrak{g}.$$

The curvature tensor is then determined by

$$\begin{array}{rcl} R : \mathfrak{m} \times \mathfrak{m} & \to & \mathfrak{gl}(\mathfrak{m}) \\ (x,y) & \mapsto & [\Lambda(x),\Lambda(y)] - \Lambda([x,y]). \end{array} \tag{5.14}$$

Finally, the Ricci tensor ϱ of g, described in terms of its components with respect to $\{u_i\}$, is given by

$$\varrho(u_i,u_j) = \sum_{r=1}^{4} R_{ri}(u_r,u_j), \qquad i,j = 1,...,4 \tag{5.15}$$

and the scalar curvature τ is the trace of ϱ.

Non-reductive homogeneous manifolds of dimension 4 were classified in [15], in terms of the corresponding non-reductive Lie algebras. The corresponding invariant pseudo-Riemannian metrics, together with their connection and curvature, were explicitly described in [12, 13]. These spaces categorized in eight classes, $A_1,\ldots,A_5,B_1,B_2,B_3$. The invariant metrics of types A_1,A_2,A_3 can be both of Lorentzian or neutral signature while the cases A_4,A_5 are always Lorentzian and cases B_1,B_2,B_3 admit the neutral signature $(2,2)$.

Theorem 5.4. *Every Lorentzian four-dimensional non-reductive homogeneous manifold is locally symmetric if and only if* $\nabla^2 R = 0$.

Proof. Let (M,g) be a Lorentzian non-reductive homogeneous four-dimensional manifold, then (M,g) is isometric to a homogeneous space G/H equipped with a Lorentzian invariant metric g, where the corresponding Lie algebras $\mathfrak{g}$ and $\mathfrak{h}$ are described in the cases $A_1,\ldots,A_5$ of [12].

We bring the details of the case A_1. The other cases can be treated in the similar way. In this case, $\mathfrak{g} = \mathcal{A}_1$ is the decomposable 5-dimensional Lie algebra $\mathfrak{sl}(2,\mathbb{R}) \oplus \mathfrak{s}(2)$, where $\mathfrak{s}(2)$ is the 2-dimensional solvable algebra. There exists a basis $\{e_1,...,e_5\}$ of $\mathcal{A}_1$, such that the non-zero products are:

$$[e_1,e_2] = 2e_2, \quad [e_1,e_3] = -2e_3, \quad [e_2,e_3] = e_1, \quad [e_4,e_5] = e_4$$

and the isotropy subalgebra is $\mathfrak{h} = \mathrm{Span}\{h_1 = e_3 + e_4\}$. So, we can take

$$\mathfrak{m} = \mathrm{Span}\{u_1 = e_1, u_2 = e_2, u_3 = e_5, u_4 = e_3 - e_4\}.$$

With respect to $\{u_i\}$, we have the following isotropy representation H_1 for h_1 and consequently the following description of invariant metric g:

$$H_1 = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & -\frac{1}{2} & 0 \end{pmatrix}, \qquad g = \begin{pmatrix} a & 0 & -\frac{a}{2} & 0 \\ 0 & b & c & a \\ -\frac{a}{2} & c & d & 0 \\ 0 & a & 0 & 0 \end{pmatrix}, \tag{5.16}$$

which is nondegenerate whenever $a(a-4d)\neq 0$ and is Lorentzian if and only if $a(a-4d)<0$. Putting $\Lambda[i]:=\Lambda(u_i)$ for all indices $i=1,...,4$, we find:

$$\Lambda[1]=\begin{pmatrix}0&0&0&0\\0&1&0&0\\0&0&0&0\\0&-\frac{b}{a}&-\frac{c}{a}&-1\end{pmatrix},\qquad \Lambda[2]=\begin{pmatrix}0&-\frac{8bd}{a(a-4d)}&\frac{c}{a}&1\\-1&0&\frac{1}{2}&0\\0&-\frac{4bc}{a(a-4d)}&0&0\\-\frac{b}{a}&\frac{4bc}{a(a-4d)}&-\frac{b}{2a}&0\end{pmatrix},\tag{5.17}$$

$$\Lambda[3]=\begin{pmatrix}0&\frac{c}{a}&0&0\\0&\frac{1}{2}&0&0\\0&0&0&0\\-\frac{c}{a}&-\frac{b}{2a}&0&-\frac{1}{2}\end{pmatrix},\qquad \Lambda[4]=0.$$

Moreover, applying (5.14) and (5.15), by setting $R_{ij}=R(u_i,u_j)$, some standard calculations give that with respect to $\{u_i\}$, the non-zero curvature components are determined as following:

$$R_{12}=\begin{pmatrix}0&\frac{b(20\,d+a)}{a(-4\,d+a)}&-\frac{c}{a}&-1\\1&0&-\frac{1}{2}&0\\0&\frac{12b}{-4\,d+a}&0&0\\\frac{4b}{a}&-\frac{12cb}{a(-4\,d+a)}&\frac{b}{a}&0\end{pmatrix},\quad R_{13}=\begin{pmatrix}0&-\frac{c}{a}&0&0\\0&0&0&0\\0&0&0&0\\\frac{c}{a}&0&-\frac{c}{2a}&0\end{pmatrix},$$

$$R_{14}=\begin{pmatrix}0&-1&0&0\\0&0&0&0\\0&0&0&0\\1&0&-\frac{1}{2}&0\end{pmatrix},\qquad R_{23}=\begin{pmatrix}0&-\frac{b(4\,d+a)}{2a(-4\,d+a)}&-\frac{c}{2a}&-\frac{1}{2}\\\frac{1}{2}&-\frac{c}{a}&-\frac{1}{4}&0\\0&-\frac{2b}{-4\,d+a}&0&0\\-\frac{b}{a}&\frac{cb(-4\,d+3\,a)}{a^2(-4\,d+a)}&\frac{c^2}{a^2}&\frac{c}{a}\end{pmatrix},$$

$$R_{24}=\begin{pmatrix}0&0&0&0\\0&-1&0&0\\0&0&0&0\\0&\frac{b}{a}&\frac{c}{a}&1\end{pmatrix},\qquad R_{34}=\begin{pmatrix}0&\frac{1}{2}&0&0\\0&0&0&0\\0&0&0&0\\-\frac{1}{2}&0&\frac{1}{4}&0\end{pmatrix},$$

and the Ricci tensor ϱ is determined by

$$\varrho=\begin{pmatrix}-2&0&1&0\\0&\frac{2b(a+12d)}{a(a-4d)}&-\frac{2c}{a}&-2\\1&-\frac{2c}{a}&-\frac{1}{2}&0\\0&-2&0&0\end{pmatrix}.$$

By using description of the curvature tensor and Levi-Civita connection, we set $\nabla R=0$ and have, the homogeneous spaces G/H is locally symmetric if and only if $b=0$. Also, $\nabla^2 R=0$ if and only if $b=0$, so we conclude that the homogeneous spaces of type A_1 are locally symmetric if and only if $\nabla^2 R=0$. Similar arguments will be applied for the other Lorentzian cases and this finishes the proof. □

The following remark is the direct conclusion of the Theorems 5.3 and 5.4.

Remark 5.5. *Let (M,g) be a homogeneous Lorentzian four-dimensional manifold, then (M,g) is not a two-symmetric space.*

Classification of four-dimensional pseudo-Riemannian homogeneous spaces with non-trivial isotropy has been studied in [19] in order to find the local classification of four-dimensional Einstein-Maxwell homogeneous spaces with an invariant pseudo-Riemannian metric of arbitrary signature. The presented list is a good reference to study pseudo-Riemannian homogeneous four-manifolds. We apply the mentioned classification to find an example of a pseudo-Riemannian two-symmetric homogeneous four-manifold, equipped with an invariant metric of *neutral* signature.

Example 5.6. *Let $M = G/H$ be a homogeneous four-dimensional manifold of type $1.4^1 : 9$ of the Komrakov's list* [19]*. In this case, the Lie algebras $\mathfrak{g}$ and $\mathfrak{h}$ are described as following:*

$$\mathfrak{g} = \left\{ \begin{pmatrix} x & 0 & 0 & 0 \\ 0 & 0 & \lambda x & 0 \\ 0 & x & x & 0 \\ 0 & 0 & x & -\mu x \end{pmatrix} \middle| \, x \in \mathbb{R}, \lambda, \mu \in \mathbb{R} \right\} \leftthreetimes (\mathfrak{n}_3 \times \mathbb{R}), \quad \mathfrak{h} = \langle p \rangle,$$

where $\mathfrak{n}_3 = \langle h, p, q \rangle$ with the only non-zero bracket $[p, q] = h$. If $\mathfrak{g} = \operatorname{span}\{u_1, u_2, u_3, u_4\}$ and $\mathfrak{h} = \operatorname{span}\{h_1\}$, the table of Lie brackets is:

$[\ ,\ ]$	h_1	u_1	u_2	u_3	u_4
h_1	0	0	u_1	u_2	0
u_1	0	0	0	u_1	0
u_2	$-u_1$	0	0	$\lambda h_1 + u_2 + u_4$	0
u_3	$-u_2$	$-u_1$	$-\lambda h_1 - u_2 - u_4$	0	μu_4
u_4	0	0	0	$-\mu u_4$	0

The invariant metric will be calculated as following

$$g = \begin{pmatrix} 0 & 0 & -a & 0 \\ 0 & a & 0 & 0 \\ -a & 0 & b & c \\ 0 & 0 & c & d \end{pmatrix},$$

for arbitrary real constants $a \neq 0, b, c, d$. This metric admits both Lorentzian and neutral signatures while for $d = -9a$, g is of neutral signature. We also set $\mu = -\frac{5}{2}$ and $\lambda = \frac{3}{4}$. Keeping in mind $\Lambda[i] = \Lambda(u_i)$ for all indices $i = 1, \ldots, 4$, the components of the Levi-Civita connection are:

$$\Lambda[1] = 0, \qquad \Lambda[2] = \begin{pmatrix} 0 & 1 & \frac{c}{2a} & -\frac{9}{2} \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & 0 \end{pmatrix},$$

$$\Lambda[3] = \begin{pmatrix} -1 & \frac{c}{2a} & \frac{6\,ba - c^2}{6a^2} & \frac{5c}{2a} \\ 0 & 0 & \frac{c}{a} & -\frac{9}{2} \\ 0 & 0 & 1 & 0 \\ 0 & -\frac{1}{2} & -\frac{c}{6a} & 0 \end{pmatrix}, \quad \Lambda[4] = \begin{pmatrix} 0 & -\frac{9}{2} & \frac{5c}{2a} & -\frac{45}{2} \\ 0 & 0 & -\frac{9}{2} & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{5}{2} & 0 \end{pmatrix},$$

also, by using Equation (5.14)*, if set* $R_{ij} = R(u_i, u_j)$*, the non-zero components of the curvature tensor are*

$$R_{23} = \begin{pmatrix} 0 & 6 & -\frac{2c}{a} & 18 \\ 0 & 0 & 6 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -2 & 0 \end{pmatrix}, \quad R_{34} = \begin{pmatrix} 0 & -18 & \frac{6c}{a} & -54 \\ 0 & 0 & -18 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 6 & 0 \end{pmatrix}.$$

The space is Ricci flat and the only non-zero covariant derivative of the curvature tensor is in the direction of u_3*. We set* $(\Lambda[k]R)_{ij} = (\Lambda(u_k)R)(u_i, u_j)$*, the non-zero components are*

$$(\Lambda[3]R)_{23} = \begin{pmatrix} 0 & 6 & -\frac{2c}{a} & 18 \\ 0 & 0 & 6 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -2 & 0 \end{pmatrix}, \quad (\Lambda[3]R)_{34} = \begin{pmatrix} 0 & -18 & \frac{6c}{a} & -54 \\ 0 & 0 & -18 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 6 & 0 \end{pmatrix}.$$

Clearly, $(M = G/H, g)$ *is never locally symmetric but by straight forward calculations we get* $\nabla^2 R = 0$ *which shows that the spaces is two-symmetric.*

References

[1] E. Abbena, S. Garbiero and L. Vanhecke, *Einstein-like metrics on three-dimensional Riemannian homogeneous manifolds*, Simon Stevin Quart. J. Pure Appl. Math. **66** (1992), 173-182.

[2] E. Abbena and S. Garbiero, *Curvature forms and Einstein-like metrics on Sasakian manifolds*, Math. J. Okayama Univ. **34** (1992), 241-248.

[3] D. V. Alekseevsky, A. S. Galaev, *Two-symmetric Lorentzian manifolds*, Geom. Phys. **61** (2011), 2331-2340.

[4] W. Ambrose, I. M. Singer, *On homogeneous Riemannian manifolds*, Duke Math. J. 25 (1958) 647-669.

[5] A. Arvanitogeorgos, *An Introduction to Lie Groups and the Geometry of Homogeneous Spaces*, Student Mathematical Library, V. 22.

[6] O. F. Blanco, M. Sánchez, J. M. Senovilla, *Complete classifcation of second-order symmetric space-times*, J. Phys. Conf. Ser. **229** (2010), 012021. 5 pp.

[7] E. Boeckx, *Einstein-like semi-symmetric spaces*, Arch. Math. (Brno) **16** (1997), 789-800.

[8] P. Bueken and L. Vanhecke, *Three- and four-dimensional Einstein-like manifolds and homogeneity*, Geom. Dedicata **75** (1999), 123-136.

[9] G. Calvaruso, *Homogeneous structures on three-dimensional Lorentzian manifolds*,J. Geom. Phys. 57 (2007), 1279-1291. Addendum: J. Geom. Phys. 58 2008, 291-292.

[10] G. Calvaruso, L. Vanhecke, *Special ball-homogeneous spaces*, Z. Anal. Anwendungen **16** (1997), 789-800.

[11] G. Calvaruso, *Einstein-like and conformally flat contact metric three-manifolds*, Balkan J. Geom. **5**(2) (2000), 17-36.

[12] G. Calvaruso, A. Fino, *Ricci solitons and geometry of four-dimensional non-reductive homogeneous spaces*, Canad. J. Math., **64** (2012), 778-804.

[13] G. Calvaruso, A. Zaeim, *Geometric structures over non-reductive homogeneous* 4*-spaces*, Adv. Geom., to appear.

[14] H. D. Cao, *Recent progress on Ricci solitons*, Recent advances in geometric analysis, 1–38, Adv. Lect. Math. (ALM), **11**, Int. Press, Somerville, MA, 2010.

[15] M. E. Fels and A.G. Renner, *Non-reductive homogeneous pseudo-Riemannian manifolds of dimension four*, Canad. J. Math. **58** (2006), 282–311.

[16] P. M. Gadea, J.A. Oubina, *Homogeneous pseudo-Riemannian structures and homogeneous almost para-Hermitian structures*, Houston J. Math. 18 (3) (1992) 449-465.

[17] A. Gray, *Einstein-like manifolds whcih are not Einstein*, Geom. Dedicata **7** (1978), 259-280.

[18] V. R. Kaigorodov, *Structure of the curvature of spacetime*, J. Sov. Laser Res. (2) **28** (1985), 256273.
[19] B. Komrakov Jnr., *Einstein-maxwell equation on four-dimensional homogeneous spaces*, Lobachevskii J. Math., **8** (2001), 33–165.
[20] V. Patrangenaru, Locally homogeneous pseudo-Riemannian manifolds, J. Geom. Phys. 17 (1995) 59-72.
[21] F. Podestá and A. Spiro, *Four-dimensional Einstein-like manifolds and curvature homogeneity*, Geom. Dedicata **54** (1995), 225-243.
[22] J. M. Senovilla, *Second-order symmetric Lorentzian manifolds. I. Characterization and general results*, Classical Quantum Gravity (24) **25** (2008), 245011. 25 pp.
[23] S. Tanno, *Curvature tensors and covariant derivatives*, Ann. Mat. Pura Appl. (4) **96** (1972), 233–241.
[24] F. Tricerri, L. Vanhecke, *Homogeneous structures on Riemannian manifolds*, in: London Math. Soc. Lect. Notes, vol. 83, Cambridge Univ. Press, 1983.
[25] J. A. Wolf, *Homogeneous manifolds of constant curvature, Comment.* Math. Helv. 36 (1961), 112-147.

Department of Mathematics, Payame noor University, P.O. Box 19395-3697, Tehran, Iran.
E-mail address: Amirhesam Zaeim: zaeim@pnu.ac.ir, Mohammad Chaichi: chaichi@pnu.ac.ir, Yadollah Keshavarzi: y.keshavarzi@pnu.ac.ir